

Table One - Pest Plants of North Point Project Area

| Name | Location | Action |
|---|--|---|
| Lantana, <i>Lantana camara</i> | Found generally throughout the whole area except in mown campground. Thickest infestations on the western side along the Honeymoon Bay (HB) track and on slopes of North Point. | Small to medium bushes removed by hand. Large bushes – base of trunk drilled and poisoned with 50% glysohate solution. Dense areas where no native plants were growing underneath, spot foliar sprayed with 10% glysohate. |
| Umbrella tree, <i>Schefflera actinophylla</i> | Mostly large individual plants scattered in the bush in the southern and eastern areas. | Large trees - base of the trunk drilled and poisoned as with Lantana. Small trees and seedlings removed by hand. |
| Easter cassia, <i>Senna pendula</i> | Several mature trees and seedlings behind fore dunes on north west side of campground. | Base of trunk drilled and poisoned with 50% glysohate. Ripe seed pods bagged. Seedlings removed by hand. |
| Prickly Pear, <i>Opuntia stricta</i> | Scattered plants found mainly around North Point and in bush areas around the campground. | Plants dug up by hand and all material bagged. |
| Asparagus fern, <i>Asparagus aethiopicus</i> | Several large plants and seedlings in the bush to the north, east, and south of the open camping areas. | Small seedlings removed by hand. Large bushes sprayed with 50% glysohate. |
| Pink periwinkle, <i>Catharanthus rosea</i> | Found generally throughout the site especially in any disturbed or open areas of bush | Mature plants and seedlings removed by hand. |
| Mossman river grass, <i>Cenchrus echinatus</i> | Thick patches in the mown areas and around facilities. Also along edges of roads and tracks. Some found in the bush adjacent to the campground | Dense patches and plants growing along road edges sprayed with 1% glysohate. Seeding plants in the campground, along tracks and in the bush removed by hand and seed heads bagged. |

| | | |
|---|--|--|
| Nightshade, <i>Solanum nigrum</i> | Ubiquitous in campground areas and perimeters | Plants and seedlings removed by hand. Mature fruit bagged |
| Dandelion, <i>Sonchus oleraceus</i> | In campground especially in and around the nursery area, and along roadsides | Plants removed by hand and seed heads bagged. Seedlings emerging in grassy areas sprayed with 1% 'Brush-Off'. |
| Khaki (bindi) weed, <i>Alternanthera pungens</i> | In open campground areas. | Plants removed by hand. Seedlings emerging in grassy areas sprayed with 1% 'Brush-Off'. |
| Broad-leaved pepper tree, <i>Schinus terebinthifolius</i> | A couple of trees in the bush at the northern end of the campground. | Trees drilled and poisoned with 50% glysophate. |
| Cobblers pegs, <i>Bidens pilulosa</i> | Ubiquitous in campground areas, perimeters, and along the edges of roads and tracks. | Plants removed by hand and seed heads bagged. Seedlings sprayed with 1% 'Brush-Off'. |
| Stinking passion vine, <i>Passiflora foetida</i> | Widespread behind fore dunes and along HB track. | Vines removed by hand. |
| Common passionfruit vine, <i>Passiflora edulis</i> | Couple of large fruiting plants found along HB track and in the adjacent bush | Vines removed by hand and ripe fruit bagged. |
| Beach primrose, <i>Oenothera drummondii</i> | In disturbed areas in mown campground and on foredunes. | Plants removed by hand. Sprayed with 1% 'Brush-Off' in grassy areas. |
| Cudweed <i>Gamochaeta sp</i> | In campground open areas and along tracks and roads | Plants removed by hand and seed heads bagged |
| Fleabane <i>Conyza sumatranum</i> | as above | as above |
| Groundsel <i>Baccharis halimifolia</i> | Two small bushes located, one at the northern end, and one along HB track. | Plants removed by hand. |
| Yucca <i>Yucca elephantipes</i> | One patch of scattered plants located in the bush north of the nursery. | Large plants slashed and poisoned with 10% glysophate. Seedlings removed by hand. |
| Rattlepod <i>Crotolaria pallida</i> | In campground and car parking areas | Plants removed by hand and seed heads bagged |

| | | |
|--|---|---|
| Live leaf <i>Bryophyllum sp</i> | One small patch found in a campsite at the southern end. | Plants removed by hand and all material bagged. |
| Rose-scented geranium <i>Pelargonium sp</i> | as above | as above |
| Thick head <i>Crassocephalum crepidiodes</i> | Ubiquitous in campground areas and perimeters | Plants removed by hand and seed heads bagged. |
| Wandering jew <i>Tradescantia zebrina</i> | One medium infestation behind the foredunes at the northern end | Plants removed by hand and all material bagged. |
| Pothos devil's ivy <i>Scindopsus aureus</i> | as above | as above. |
| Bauhinia <i>Bauhinia sp</i> | One specimen on edge of campground | Drilled and poisoned with glysophate. |
| Grevillea <i>Grevillea venusta</i> | One tree adjacent to campground | no action. |
| Fig tree <i>Ficus macrocarpa</i> | One large tree in campground adjacent to nursery | no action |
| Grasses; Green panic <i>Panicum maximum</i> , African lovegrass <i>Eragrostis curvula</i> , Signal grass <i>Brachiaria decumbens</i> | Along edges of main road adjacent to the campground. | no action |